ELSEVIER

Journal of Statistical Planning and Inference 56 (1996) 325–326

journal of statistical planning and inference

Author Index Volume 56 (1996)*

Adams, P., E.J. Billington and D.E. Bryant, The spectrum for lambda-fold Steiner pentagon systems (1) 3–15
Assmus, Jr., E.F., On Berman's characterization of the Reed-Muller codes (1) 17–22
Baker, R.D., G.L. Ebert, J. Hemmeter and A. Woldar, Maximal cliques in the Paley graph of square order (1) 33–38
Banerjee, S. and S. Kageyama, Non-isomorphic solutions of group divisible designs with $k = 3$ and $\lambda_1 = 0$ (1) 23–32
Bierbrauer, J., Construction of orthogonal arrays (1) 39–47
Billington, E.J., *see* P. Adams (1) 3–15
Bloemen, I., J.A., Thas, H. Van Maldeghem, Elation generalized quadrangles of order (p, t), p prime, are classical (1) 49–55
Bryant, D.E., *see* P. Adams (1) 3–15
Bryant, D.E. and C.C. Lindner, 2-perfect directed m-cycle systems can be equationally defined for $m = 3$, 4, and 5 only (1) 57–63
Camina, A.R. and A.A. Miller, Line-closed subsets of Steiner triple systems and classical linear spaces (1) 65–77
Charpin, P., Self-dual codes which are principal ideals of the group algebra $\mathbb{F}_2[\{\mathbb{F}_{2^m}, +\}]$ (1) 79–92
Colbourn, C.J., Some direct constructions for incomplete transversal designs (1) 93–104
Ebert, G.L., *see* R.D. Baker (1) 33–38
Fuji-Hara, R., S. Kuriki and M. Miyake, Cyclic orthogonal and balanced arrays (2) 171–180
Fujii, Y., *see* S. Yamamoto (2) 269–287
Hamada, N. and T. Helleseth, The uniqueness of [87, 5, 57; 3]-codes and the nonexistence of [258, 6, 171; 3]-codes (1) 105–127
Hamada, N. and T. Helleseth, A characterization of some $\{3v_2 + v_3, 3v_1 + v_2; 3, 3\}$-minihypers and some [15, 4, 9; 3]-codes with $B_2 = 0$ (1) 129–146
Hamada, N. and T. Maekawa, A characterization of some $\{3v_1 + v_3, 3v_0 + v_2; 3, 3\}$-minihypers and its applications to error-correcting codes (1) 147–169
Hedayat, A.S., J. Stufken and G. Su, On difference schemes and orthogonal arrays of strength t (2) 307–324
Helleseth, T., *see* N. Hamada (1) 105–127
Helleseth, T., *see* N. Hamada (1) 129–146
Hemmeter, J., *see* R.D. Baker (1) 33–38
Hyodo, Y., *see* S. Yamamoto (2) 269–287
Kageyama, S., *see* S. Banerjee (1) 23–32
Key, J.D., Extendable Steiner designs from finite geometries (2) 181–186
Kuriki, S., *see* R. Fuji-Hara (2) 171–180
Lam, C. and V.D. Tonchev, Classification of affine resolvable 2-(27, 9, 4) designs (2) 187–202
Lindner, C.C., *see* D.E. Bryant (1) 57–63
Maekawa, T., *see* N. Hamada (1) 147–169
Mandrell. R. and J. Yucas, A geometric construction of the Steiner system $S(5, 8, 24)$ (2) 223–228
Mathon, R. and C.E. Praeger, On block-transitive designs with affine automorphism group (2) 203–221
Mesner, D.M., *see* T. van Trung (2) 257–268

*The issue number is given in front of page numbers

Elsevier Science B.V.

Miller, A.A., *see* A.R. Camina (1) 65–77
Miyake, M., *see* R. Fuji-Hara (2) 171–180
Moorhouse, G.E., Some p-ranks related to Hermitian varieties (2) 229–241
Praeger, C.E., *see* R. Mathon (2) 203–221
Shirakura, T. and W.-P. Tong, Weighted A-optimality for fractional 2^m factorial designs of resolution V (2) 243–256
Sloane, N.J.A. and J. Stufken, A linear programming bound for orthogonal arrays with mixed levels (2) 295–305
Stufken, J., *see* N.J.A. Sloane (2) 295–305
Stufken, J., *see* A.S. Hedayat (2) 307–324
Su, G., *see* A.S. Hedayat (2) 307–324
Thas, J.A., *see* I. Bloemen (1) 49–55
Tonchev, V.D., *see* C. Lam (2) 187–202
Tong, W.-P., *see* T. Shirakura (2) 243–256
Van Maldeghem, H., *see* I. Bloemen (1) 49–55
van Trung, T., Q.-r. Wu and D.M. Mesner, High order intersection numbers of t-designs (2) 257–268
Woldar, A., *see* R.D. Baker (1) 33–38
Wu, Q.-r., *see* T. van Trung (2) 257–268
Yamamoto, S., Y. Fujii, Y. Hyodo and H. Yumiba, New light on fractional 2^m factorial designs (2) 269–287
Yucas, J., *see* R. Mandrell (2) 223–228
Yumiba, H., *see* S. Yamamoto (2) 269–287
Zinoviev, V.A., On the equivalence of certain constant weight codes and combinatorial designs (2) 289–294